


— —

IYGB - M1 PAPER T - QUESTION 1

a)



- This is the standard
 $y = \frac{1}{x^2}$ turned upside


Down"

- Asymptotes

$x=0$ (y axis)

$y=0$ (x axis)

b) I)



- This is a translation
by 1 unit to the "right"


- Asymptotes

$x=1$

$y=0$ (x axis)

II)

$$f'(x) = \frac{d}{dx} \left(-\frac{1}{x^2} \right) = \frac{2}{x^3}$$



- This in the absence of scale looks like $\frac{a}{x}$

- Asymptotes

$x=0$ (y axis)

$y=0$ (x axis)

-1-

IYGB-MPI PAPER 0 - QUESTION 2

FORMING THE EQUATION & TIDY TO A 3 TERM QUADRATIC

$$\Rightarrow px^2 + 4x(p+3) + 5p = -19$$

$$\Rightarrow \underbrace{px^2}_a + \underbrace{4(p+3)x}_b + \underbrace{(5p+19)}_c = 0$$

TWO DISTINCT ROOTS IMPLY $b^2 - 4ac > 0$

$$\Rightarrow [4(p+3)]^2 - 4 \times p \times (5p+19) > 0$$

$$\Rightarrow 16(p+3)^2 - 4p(5p+19) > 0$$

$$\Rightarrow 4(p+3)^2 - p(5p+19) > 0$$

$$\Rightarrow 4(p^2 + 6p + 9) - 5p^2 - 19p > 0$$


$$\Rightarrow 4p^2 + 24p + 36 - 5p^2 - 19p > 0$$

$$\Rightarrow -p^2 + 5p + 36 > 0$$

$$\Rightarrow p^2 - 5p - 36 < 0$$

$$\Rightarrow (p-9)(p+4) < 0$$

CRITICAL VALUES ARE 9 & -4



$$\text{so } -4 < p < 9$$

BUT $p \neq 0$, OTHERWISE NO QUADRATIC

$$-4 < p < 0 \cup 0 < p < 9$$

- 1 -

IYGB - MPI PAPER 0 - QUESTION 3

GETTING RID OF THE DENOMINATORS

$$\Rightarrow \frac{2 + \cos 2x}{3 + \sin^2 2x} = \frac{2}{5}$$

$$\Rightarrow 10 + 5\cos 2x = 6 + 2\sin^2 2x$$

USING $\cos^2 2x + \sin^2 2x \equiv 1$

$$\Rightarrow 10 + 5\cos 2x = 6 + 2(1 - \cos^2 2x)$$

$$\Rightarrow 10 + 5\cos 2x = 6 + 2 - 2\cos^2 2x$$

$$\Rightarrow 2\cos^2 2x + 5\cos 2x + 2 = 0$$

$$\Rightarrow (2\cos 2x + 1)(\cos 2x + 2) = 0$$

$$\Rightarrow \cos 2x = \begin{cases} -2 & -1 \leq \cos 2x \leq 1 \\ -\frac{1}{2} \end{cases}$$


PROCEED WITH SOLUTION

$$\arccos(-\frac{1}{2}) = 120^\circ$$

$$\begin{cases} 2x = 120^\circ \pm 360n & n=0,1,2,3,\dots \\ 2x = 240^\circ \pm 360n \end{cases}$$

$$\begin{cases} x = 60^\circ \pm 180n \\ x = 120^\circ \pm 180n \end{cases}$$

$x = 60^\circ, 240^\circ, 120^\circ, 300^\circ$



IYGB - MPI PAPER U - QUESTION 4

a) $x-2$ IS A FACTOR OF $f(x)$; $2x+1$ IS A FACTOR OF $f(x)$

$$f(2) = 0$$

$$2 \times 2^3 - 9 \times 2^2 + px \times 2 + q = 0$$

$$16 - 36 + 2p + q = 0$$

$$2p + q = 20$$

$$q = 20 - 2p$$

$$f(-\frac{1}{2}) = 0$$

$$2(-\frac{1}{2})^3 - 9 \times (-\frac{1}{2})^2 + p(-\frac{1}{2}) + q = 0$$

$$-\frac{1}{4} - \frac{9}{4} - \frac{1}{2}p + q = 0$$

$$-\frac{5}{2} - \frac{1}{2}p + q = 0$$

$$q = \frac{1}{2}p + \frac{5}{2}$$

$$\begin{aligned} & \rightarrow 20 - 2p = \frac{1}{2}p + \frac{5}{2} \\ & 40 - 4p = p + 5 \end{aligned}$$

$$35 = 5p$$

$$p = 7$$

$$q = 6$$

b) $2\sqrt{y} + \frac{7}{\sqrt{y}} = 9 - \frac{6}{y}$

LET $\alpha = \sqrt{y}$

$$\Rightarrow 2\alpha + \frac{7}{\alpha} = 9 - \frac{6}{\alpha^2}$$

$$\Rightarrow 2\alpha^3 + 7\alpha = 9\alpha^2 - 6$$

$$\Rightarrow 2\alpha^3 - 9\alpha^2 + 7\alpha + 6 = 0$$

THIS IS THE CUBIC OF PART (a) — FACTORIZE BY INSPECTION

$$\Rightarrow (2x+1)(x-2)(x-3) = 0$$

$$\Rightarrow x = \begin{cases} 3 \\ 2 \\ -\frac{1}{2} \end{cases}$$

-2 -

IYGB - MPI PAPER U - QUESTION 4

$$\Rightarrow \sqrt{y} = \sqrt{2} \quad \sqrt{3}$$

~~$\frac{1}{2}$~~

$$\Rightarrow y = 4 \quad 9$$

 |||

- / -

IYGB - MPI PAPER U - QUESTION 5

METHOD A - BY COORDINATE GEOMETRY


- GRADIENT OF THE TANGENT IS 2
- GRADIENT CT MUST BE $-\frac{1}{2}$
- EQUATION OF LINE THROUGH C & T

$$y - (-3) = -\frac{1}{2}(x - 2)$$

$$y + 3 = -\frac{1}{2}(x - 2)$$

$$2y + 6 = -x + 2$$

$$2y + x + 4 = 0$$



- SOLVING SIMULTANEOUSLY WITH $y = 2x - 3$

$$2(2x - 3) + x + 4 = 0$$

$$5x - 2 = 0$$

$$x = \frac{2}{5}$$

$$\text{AND } y = 2\left(\frac{2}{5}\right) - 3 = \frac{4}{5} - 3 = -\frac{11}{5} \quad \therefore \left(\frac{2}{5}, -\frac{11}{5}\right)$$

- DISTANCE CT FINALLY, $C(2, -3)$ & $T\left(\frac{2}{5}, -\frac{11}{5}\right)$

$$d = \sqrt{(y_2 - y_1)^2 + (x_2 - x_1)^2}$$

$$|CT| = \sqrt{\left(-3 + \frac{11}{5}\right)^2 + \left(\frac{2}{5} - 2\right)^2}$$

$$r = \sqrt{\left(-\frac{4}{5}\right)^2 + \left(-\frac{8}{5}\right)^2} = \sqrt{\frac{16}{25} + \frac{64}{25}} = \sqrt{\frac{80}{25}} = \frac{4}{5}\sqrt{5}$$

METHOD B - USING DISCRIMINANTS

- LET THE CIRCLE HAVE EQUATION

$$(x - 2)^2 + (y + 3)^2 = r^2$$

-2-

IYGB - MPI PAGE 6 - QUESTION 5

- SOLVING SIMULTANEOUSLY WITH $y = 2x - 3$ TO "FIND" T

$$\Rightarrow (x-2)^2 + (2x-3+3)^2 = r^2$$

$$\Rightarrow (x-2)^2 + (2x)^2 = r^2$$

$$\Rightarrow x^2 - 4x + 4 + 4x^2 = r^2$$

$$\Rightarrow 5x^2 - 4x + (4 - r^2) = 0$$

- THIS EQUATION MUST PRODUCE REPEATED ROOTS AS THE POINT T IS A POINT OF TANGENCY

$$b^2 - 4ac = 0 \Rightarrow (-4)^2 - 4 \times 5 \times (4 - r^2) = 0$$

$$\Rightarrow 16 - 20(4 - r^2) = 0$$

$$\Rightarrow 16 - 80 + 20r^2 = 0$$

$$\Rightarrow 20r^2 = 64$$

$$\Rightarrow r^2 = \frac{64}{20} = \frac{64}{100} \times 5$$

$$\Rightarrow r = \underline{\underline{\frac{4}{5}\sqrt{5}}} \quad \text{AS BEFORE}$$

METHOD C - BY MINIMIZATION (COMPLETING THE SQUARE)

- CONSIDER A POINT ON THE LINE $y = 2x - 3$, i.e. $(x, 2x-3)$

- THE DISTANCE FROM $(x, 2x-3)$ TO THE CENTER $(2, -3)$ IS GIVEN BY

$$\Rightarrow d = \sqrt{(x-2)^2 + [2x-3 - (-3)]^2}$$

$$\Rightarrow d = \sqrt{(x-2)^2 + 4x^2}$$

$$\Rightarrow d^2 = x^2 - 4x + 4 + 4x^2$$

-3-

LYGB - MFL PAPER U - QUESTION 5

$$\Rightarrow d^2 = 5x^2 - 4x + 4$$

$$\Rightarrow d^2 = 5 \left[x^2 - \frac{4}{5}x + \frac{4}{5} \right]$$

$$\Rightarrow d^2 = 5 \left[\left(x - \frac{2}{5} \right)^2 - \frac{4}{25} + \frac{4}{5} \right]$$

$$\Rightarrow d^2 = 5 \left(x - \frac{2}{5} \right)^2 - \frac{4}{5} + 4$$

$$\Rightarrow d^2 = 5 \left(x - \frac{2}{5} \right)^2 + \frac{16}{5}$$

\therefore MINIMUM VALUE OF d^2 IS $\frac{16}{5}$ (occurs at $x = \frac{2}{5}$)

$$\therefore d_{\text{MIN}} = r = \sqrt{\frac{16}{5}} = \sqrt{\frac{16 \times 5}{25}} = \frac{4}{5}\sqrt{5}$$

-1-

IYGB-MPI PAPER T - QUESTION 6

C: $y = ax^{\frac{3}{2}} + bx^{\frac{1}{2}}, x > 0$ • L: $y = 8x - 32$

• "L IS A TANGENT TO THE WAVE AT $x=4$ "

• $x=4$

$$y = 8 \times 4 - 32$$

$$y = 0$$

• GRADIENT OF L IS 8

• $y = ax^{\frac{3}{2}} + bx^{\frac{1}{2}}$

$$\frac{dy}{dx} = \frac{3}{2}ax^{\frac{1}{2}} + \frac{1}{2}bx^{-\frac{1}{2}}$$

• WHEN $x=4, \frac{dy}{dx}=8$

$$\Rightarrow 8 = \frac{3}{2}a \times 4^{\frac{1}{2}} + \frac{1}{2}b \times 4^{-\frac{1}{2}}$$

$$\Rightarrow 8 = 3a + \frac{1}{4}b$$

$$32 = 12a - b$$

• WHEN $x=0, y=0$

$$\Rightarrow 0 = a \times 4^{\frac{3}{2}} + b \times 4^{\frac{1}{2}}$$

$$\Rightarrow 0 = 8a - 2b$$

$$\Rightarrow 2b = 8a$$

$$\Rightarrow b = 4a$$

• SOLVING THE EQUATIONS METHODS

$$\Rightarrow 32 = 12a - (4a)$$

$$\Rightarrow 32 = 8a$$

$$\Rightarrow a = 4$$

a
b = 16



IYGB - MPP PAPER U - QUESTION 7

LOOKING AT THE DIAGRAM

$$\Rightarrow |AB|^2 + |BC|^2 = |AC|^2$$

$$\Rightarrow 9^2 + 21^2 = |AC|^2$$

$$\Rightarrow |AC|^2 = 522$$

$$\Rightarrow |AC| = \sqrt{522}$$

BY THE SINE RULE ON $\triangle ACD$

$$\frac{\sin \theta}{24} = \frac{\sin 32^\circ}{\sqrt{522}} \Rightarrow \sin \theta = \frac{24 \sin 32^\circ}{\sqrt{522}}$$

$$\Rightarrow \sin \theta = 0.55665\dots$$

$$\Rightarrow \theta = 33.8247^\circ \quad \leftarrow \text{NOT OBTUSE AS } \sqrt{522} \approx 22.8 \text{ WHICH IS COMPARABLE WITH } 24$$

FINALLY FIND ϕ & THE AREA OF $\triangle ACD$

$$\phi = 180 - (32 + 33.8247\dots)$$

$$\phi = 114.1753\dots^\circ$$

$$\begin{aligned}\text{AREA OF } \triangle ACD &= \frac{1}{2} |AC| |CD| \sin \phi \\ &= \frac{1}{2} \sqrt{522} \times 24 \times \sin(114.1753\dots^\circ) \\ &= 250.122\dots \text{ cm}^2\end{aligned}$$

NEXT THE AREA OF $\triangle ABC$


$$\text{AREA} = \frac{1}{2} |AB| |BC| = \frac{1}{2} \times 9 \times 21 = 94.5 \text{ cm}^2$$

$$\therefore \text{REQUIRED AREA} = 250.122\dots + 94.5$$

$$= 344.622\dots$$

$$\simeq 345 \text{ cm}^2$$

3 s.f.



- -

IIT-JEE-NET PAPER U - QUESTION 8

PROCEED AS FOLLOWS

$$\Rightarrow \frac{\log_4 x^2}{5 + \log_4 x^2} + (\log_4 x)^2 = 0$$

$$\Rightarrow \frac{2\log_4 x}{5 + 2\log_4 x} + (\log_4 x)^2 = 0$$

$$\Rightarrow \frac{2y}{5 + 2y} + y^2 = 0$$

$$\Rightarrow 2y + y^2(5 + 2y) = 0 (5 + 2y)$$

$$\Rightarrow 2y + 5y^2 + 2y^3 = 0$$

$$\Rightarrow 2y^3 + 5y^2 + 2y = 0$$

$$\Rightarrow y(2y^2 + 5y + 2) = 0$$

$$\Rightarrow y(2y + 1)(y + 2) = 0$$

$$\Rightarrow y = \begin{cases} 0 \\ -\frac{1}{2} \\ -2 \end{cases}$$

i.e.

$$\log_4 x = \begin{cases} 0 \\ -\frac{1}{2} \\ -2 \end{cases}$$

INVERTING BY INSPECTION

$$x = \begin{cases} 4^0 \\ 4^{-\frac{1}{2}} \\ 4^{-2} \end{cases}$$

i.e.

$$x = \begin{cases} 1 \\ \frac{1}{2} \\ \frac{1}{16} \end{cases}$$

- -

IYGB - MPI PAPER 0 - QUESTION 9

A(-1,4) • B(2,3) • C(8,1)


a) FIND THE VECTORS \vec{AB} & \vec{BC}

$$\vec{AB} = \underline{b} - \underline{a} = \begin{pmatrix} 2 \\ 3 \end{pmatrix} - \begin{pmatrix} -1 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 \\ -1 \end{pmatrix}$$

$$\vec{BC} = \underline{c} - \underline{b} = \begin{pmatrix} 8 \\ 1 \end{pmatrix} - \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 6 \\ -2 \end{pmatrix} = 2 \begin{pmatrix} 3 \\ -1 \end{pmatrix}$$

AS \vec{AB} & \vec{BC} ARE IN THE SAME DIRECTION & SHARE THE POINT B, A, B & C MUST BE COLLINEAR.

b)



WORKING AT THE DIAGRAM

$$\Rightarrow \vec{OD} = \vec{OB} + \vec{BD}$$

$$\Rightarrow \vec{OD} = \vec{OB} + \frac{2}{3} \vec{BC}$$

$$\Rightarrow \underline{d} = \underline{b} + \frac{2}{3}(\underline{c} - \underline{b})$$

$$\Rightarrow 3\underline{d} = 3\underline{b} + 2\underline{c} - 2\underline{b}$$

$$\Rightarrow 3\underline{d} = \underline{b} + 2\underline{c}$$

$$\Rightarrow 3\underline{d} = \begin{pmatrix} 2 \\ 3 \end{pmatrix} + 2 \begin{pmatrix} 8 \\ 1 \end{pmatrix}$$

$$\Rightarrow 3\underline{d} = \begin{pmatrix} 18 \\ 5 \end{pmatrix}$$


$$\Rightarrow \underline{d} = \begin{pmatrix} 6 \\ \frac{5}{3} \end{pmatrix}$$

$$\therefore \underline{D}\left(6, \frac{5}{3}\right)$$

-2-

IYGB - MPI PAPER 7 - QUESTION 9

b) WORKING AT THE DIAGRAM BELOW



$$\Rightarrow \vec{OP} = \vec{OA} + \vec{AP}$$

$$\Rightarrow \gamma \vec{OB} = \vec{OA} + \mu \vec{OC}$$

$$\Rightarrow \gamma \underline{b} = \underline{a} + \mu \underline{c}$$

$$\Rightarrow \gamma \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 4 \end{pmatrix} + \mu \begin{pmatrix} 8 \\ 1 \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} 2\lambda \\ 3\lambda \end{pmatrix} = \begin{pmatrix} -1 + 8\mu \\ 4 + \mu \end{pmatrix}$$

$$\Rightarrow \frac{2\lambda}{-24\lambda} = \frac{-1 + 8\mu}{-32 - 8\mu}$$

$$\Rightarrow -22\lambda = -33$$

$$\Rightarrow \gamma = \frac{3}{2}$$

Hence as $\vec{OP} = \gamma \vec{OB}$

$$\vec{OP} = \frac{3}{2} \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 3 \\ 9/2 \end{pmatrix}$$

$$\therefore \underline{P} \begin{pmatrix} 3, \frac{9}{2} \end{pmatrix}$$

- i -

IGCSE - MATH PAPER 7 - QUESTION 10

a) EXPANDING BY THE STANDARD BINOMIAL FORMULA

$$\begin{aligned}(6x-3)^8 &= \binom{8}{0}(6x)^0(-3)^8 + \binom{8}{1}(6x)^1(-3)^7 + \binom{8}{2}(6x)^2(-3)^6 + \binom{8}{3}(6x)^3(-3)^5 + \dots \\&= (1 \times 1 \times 6561) + (8 \times 6x \times (-2187)) + (28 \times 36x^2 \times 729) \\&\quad + (56 \times 216x^3 \times (-243)) + \dots \\&= \underline{\underline{6561 - 104976x + 734832x^2 - 2939328x^3}} + \dots\end{aligned}$$

b) PROCEED AS FOLLOWS

$$\frac{y+9}{3} = 6x+3$$

$$y+9 = 18x+9$$

$$18x = y$$

$$x = \frac{1}{18}y$$

USING PART (a)

$$\left[6\left(\frac{1}{18}y\right)-3\right]^8 = 6561 - \dots - 2939328\left(\frac{1}{18}y\right)^3 + \dots$$

$$\left[\frac{y+9}{3}\right]^8 = 6561 - \dots - 504y^3 + \dots$$

~~lt 504~~

c) WORK IT OUTWARDS

$$\begin{aligned}(\sqrt{2}z-1)^8 (\sqrt{2}z+1)^8 &= [(\sqrt{2}z-1)(\sqrt{2}z+1)]^8 \\&= (2z^2-1)^8 \\&= \frac{1}{3^8} \times 3^8 \times (2z^2-1)^8\end{aligned}$$

-2 -

(YGB - MPI PAPER V - QUESTION 10)

$$= \frac{1}{3^8} \times [3(2z^2 - 1)]^8$$

$$= \frac{1}{6561} [6z^2 - 3]^8$$

$$= \frac{1}{6561} \left[6561 - 104976(z^2) + 734832(z^2)^2 - 2939328(z^2)^3 + \dots \right]$$

$$= \frac{1}{6561} \left[\dots - 2939328z^6 + \dots \right]$$

$$= \dots - 448z^6 + \dots$$

1.E - 448



- -

IYGBI - MPI PAPER U - QUESTION 11.

a)

- START WITH A DIAGRAM (NOT TO SCALE)


$$\Rightarrow 3x + y = 12$$

$$\Rightarrow y = -3x + 12$$

- GRADIENT OF \overline{ABC} WILL BE $+\frac{1}{3}$

- EQUATION OF LINE \overline{ABC}

$$y - 1 = \frac{1}{3}(x + 2)$$



- SOLVING SIMULTANEOUSLY WITH $3x + y = 12$

$$\begin{aligned} y - 1 &= \frac{1}{3}(x + 2) \\ y &= 12 - 3x \end{aligned} \quad \left\{ \Rightarrow (12 - 3x) - 1 = \frac{1}{3}(x + 2) \right.$$

$$\Rightarrow 11 - 3x = \frac{1}{3}(x + 2)$$

$$\Rightarrow 33 - 9x = x + 2$$

$$\Rightarrow 31 = 10x$$

$$\Rightarrow x = 3.1$$

$$\therefore y = 12 - 3(3.1) = 12 - 9.3 = 2.7$$

Thus so far $B(3.1, 2.7)$

- NOW THE REQUIRED POINT C MUST BE SUCH SO THAT B IS THE MIDPOINT OF AC

$$\begin{array}{c} (-2) \xrightarrow{+5.1} (3.1) \xrightarrow{+5.1} (8.2) \\ \downarrow \qquad \uparrow \qquad \uparrow \\ A \qquad B \qquad C \end{array}$$

$$\therefore C(8.2, 4.4)$$

IYGB - MPI PAPER V - QUESTION II

b)

- STARTING WITH A DIAGRAM AGAIN


- GRADIENT $AP = \frac{2-1}{4+2} = \frac{1}{6}$

- THUS GRADIENT $AQ = -6$

- $|AP| = \sqrt{(2-1)^2 + (4+2)^2}$

$$|AP| = \sqrt{1+36}$$

$$|AP| = \sqrt{37}$$



- THE EQUATION OF THE LINE THROUGH A & Q IS

$$\Rightarrow y-1 = -6(x+2)$$

- THIS MUST BE SATISFIED BY Q(a,b)

$$\Rightarrow b-1 = -6(a+2)$$

- ALSO THE DISTANCE $AQ = \sqrt{37}$

$$\Rightarrow \sqrt{(a+2)^2 + (b-1)^2} = \sqrt{37}$$

$$\Rightarrow (a+2)^2 + (b-1)^2 = 37$$

$$\Rightarrow (a+2)^2 + [-6(a+2)]^2 = 37$$

$$\Rightarrow (a+2)^2 + 36(a+2)^2 = 37$$

$$\Rightarrow 37(a+2)^2 = 37$$

$$\Rightarrow (a+2)^2 = 1$$

$$\Rightarrow a+2 = \begin{cases} 1 \\ -1 \end{cases}$$

$$\Rightarrow a = \begin{cases} -1 \\ -3 \end{cases} \quad b = \begin{cases} -5 \\ 7 \end{cases}$$

~~(-1, 5)~~
~~(-3, 7)~~

$\therefore Q(-3, 7)$

-1-

YGB - MPI PAPER 75 - QUESTION 12

LOOKING AT THE FIRST EQUATION

$$\int_1^2 kx^2 + a \, dx = 11 \Rightarrow \left[\frac{1}{3}kx^3 + ax \right]_1^2 = 11$$
$$\Rightarrow \left(\frac{8}{3}k + 2a \right) - \left(\frac{1}{3}k + a \right) = 11$$
$$\Rightarrow \frac{7}{3}k + a = 11$$

LOOKING AT THE SECOND EQUATION

$$\int_1^k \frac{6}{x^2} \, dx = a \Rightarrow \left[-\frac{6}{x} \right]_1^k = a$$
$$\Rightarrow -\frac{6}{k} + 6 = a$$

SOLVING SIMULTANEOUSLY BY SUBSTITUTION

$$\Rightarrow \frac{7}{3}k + \left(-\frac{6}{k} + 6 \right) = 11$$
$$\Rightarrow \frac{7}{3}k - \frac{6}{k} + 6 = 11 \quad \rightarrow \times 3k$$
$$\Rightarrow 7k^2 - 18 + 18k = 33k$$
$$\Rightarrow 7k^2 - 15k - 18 = 0$$
$$\Rightarrow (7k+6)(k-3) = 0$$

$$k = \begin{cases} 3 \\ \cancel{-\frac{6}{7}} \end{cases}$$